



Online Appendix



NS, not significant.

FIGURE 1: Annual rainfall totals for four climate stations, (a) Spekbos, (b) Wildebeeskom, (c) Gannaleegte and (d) Nqweba Dam, within the Camdeboo National Park over the period 1987–2010.



NS, not significant.

FIGURE 2: Long-term annual maximum and minimum temperature for the study area (Graaff-Reinet) from 1992–2008.





FIGURE 3: Changes in annual S-pan evaporation (mm) values for Nqweba Dam over the period 1989–2010.

Note: This is the Online Appendix of Masubelele, M.L., Hoffman, M.T., Bond, W. & Burdett, P., 2013, 'Vegetation change (1988–2010) in Camdeboo National Park (South Africa), using fixed-point photo monitoring: The role of herbivory and climate', *Koedoe* 55(1), Art. #1127, 16 pages. <http://dx.doi.org/10.4102/koedoe.v55i1.1127>



LSU, large stock equivalent.

FIGURE 4: Herbivore density at Camdeboo National Park as shown by average stocking density (LSU/km²) for (a) 1990, (b) 1991, (c) 1992, (d) 1993, (e) 1994, (f) 1995, (g) 1996, (h) 1997, (i) 1998, (j) 1999, (k) 2000, (l) 2001, (m) 2002, (n) 2003, (o) 2004, (p) 2005, (q) 2006, (r) 2007, (s) 2008, (t) 2009 and (u) 2010.