

# AN ECOLOGICAL STUDY OF THE PLANT COMMUNITIES IN THE PROPOSED HIGHVELD NATIONAL PARK, IN THE PERI-URBAN AREA OF POTCHEFSTROOM, SOUTH AFRICA

**Authors:**

Mahlomola E. Daemane<sup>1</sup>  
Sarel S. Cilliers<sup>2</sup>  
Hugo Bezuidenhout<sup>1</sup>

**Affiliations:**

<sup>1</sup>Conservation Services  
Department, South African  
National Parks,  
South Africa

<sup>2</sup>School of Environmental  
Sciences and  
Development, North-West  
University, South Africa

**Correspondence to:**

Mahlomola Daemane

**email:**

ernestd@sanparks.org

**Postal address:**

Conservation Services  
Department, PO Box  
110040, Hadison Park,  
Kimberley 8306,  
South Africa

**Keywords:**

Braun-Blanquet;  
conservation; degradation;  
Grassland Biome;  
ordination

**Dates:**

Received: 07 May 2009  
Accepted: 11 Dec. 2009  
Published: 26 Apr. 2010

**How to cite this article:**

Daemane, M.E., Cilliers,  
S.S. & Bezuidenhout, H.,  
2010, 'An ecological study  
of the plant communities  
in the proposed Highveld  
National Park, in the  
peri-urban area of  
Potchefstroom, South  
Africa', *Koedoe* 52(1),  
Art. #708, 8 pages. DOI:  
10.4102/koedoe.v52i1.708

**This article is available at:**

<http://www.koedoe.co.za>

© 2010. The Authors.  
Licensee: OpenJournals  
Publishing. This work  
is licensed under the  
Creative Commons  
Attribution License.

**ABSTRACT**

The proposed Highveld National Park (HNP) is an area of high conservation value in South Africa, covering approximately 0.03% of the endangered Grassland Biome. The park is situated immediately adjacent to the town of Potchefstroom in the North-West Province. The objective of this study was to identify, classify, describe and map the plant communities in this park. Vegetation sampling was done by means of the Braun-Blanquet method and a total of 88 stratified random relevés were sampled. A numerical classification technique (TWINSPAN) was used and the results were refined by Braun-Blanquet procedures. The final results of the classification procedure were presented in the form of phytosociological tables and, thereafter, nine plant communities were described and mapped. A detrended correspondence analysis confirmed the presence of three structural vegetation units, namely woodland, shrubland and grassland. Differences in floristic composition in the three vegetation units were found to be influenced by environmental factors, such as surface rockiness and altitude. Incidences of harvesting trees for fuel, uncontrolled fires and overgrazing were found to have a significant effect on floristic and structural composition in the HNP. The ecological interpretation derived from this study can therefore be used as a tool for environmental planning and management of this grassland area.

**Conservation implications:** Grasslands are amongst the most threatened and least conserved vegetation biomes in South Africa, with less than 1.3% currently being conserved. The HNP has significant value for biodiversity conservation and the protection of this area will contribute to the preservation of the highly threatened Highveld vegetation types.


**INTRODUCTION**

Conservation strategies are being developed to expand the number and size of protected areas by incorporating communal lands. South Africa stands at the threshold of important and difficult socio-political decisions with respect to land restitution and redistribution (Fabricius, Koch & Magome 2001; Hoffman & Meadows 2002). New policies allow communities better access to natural resources and call for their participation in the management of protected areas, as well as facilitate the restitution of land from which they had been forcibly removed (Fabricius *et al.* 2001). Conservation agencies have, in some instances, been able to expand the size of the protected wildlife estate by entering into negotiations with local residents. This has resulted in a new category of protected areas called 'contractual parks'. In South Africa, examples of this co-management strategy include the Richtersveld National Park and the agreement with the Makuleke people in the Kruger National Park (Archer 1999; Steenkamp 1999). Although the Highveld National Park (HNP) is not regarded as a contractual park, all the stakeholders (South African National Parks, the Barolong Community, North-West Parks and Tourism Board, and the Tlokwe City Council of Potchefstroom) constitute the 'Founding Partners', with a memorandum of understanding regulating their respective rights and obligations within the park. This agreement defines the roles and responsibilities of the different partners, including the management and administration of the HNP.

The establishment of the HNP aimed at conserving a considerable area of the poorly conserved Rocky Highveld Grassveld (Bredenkamp & Van Rooyen 1996a) and Dry Sandy Highveld Grassveld (Bredenkamp & Van Rooyen 1996b) of the western Grassland Biome in South Africa; areas which have been considered to be the most scenic landscapes in the western Grassland Biome (Mucina & Rutherford 2006). According to Fairbanks *et al.* (2000), almost 30% of the Grassland Biome of South Africa has been permanently transformed, primarily as a result of cultivation (23%), plantation forestry (4%), urbanisation (2%) and mining (1%). A further 7% has been severely degraded by erosion and agricultural development. In order to maintain biodiversity, it is important to conserve elements such as vegetation and soil, because their loss enhances erosion and reduces the productive value of the land (Hoffman & Todd 2000; O'Brien 2004). Urbanisation is a major contributor to the loss of natural areas in certain parts of the Grassland Biome (Cilliers, Müller & Drewes 2004), particularly in the Potchefstroom area. Although urbanisation in some provinces has been lower than South Africa's national figure of 53%, a rapid increase is expected in the future (Cilliers *et al.* 2004), as a result of the higher levels of poverty and fewer job opportunities in rural areas that have been caused by continued droughts, specifically in the North-West Province. The identification and understanding of plant communities, as well as their associated habitats, are of fundamental importance in the formulation and execution of conservation strategies for the HNP.

**STUDY AREA****Location**

The study focused on the area west of Potchefstroom (Figure 1), between 26°39'36" S – 26°46'12" S and 26°52'12" E – 27°04'12" E. The area for park development lies between the N12, running from Potchefstroom to Klerksdorp in the south and Eleazer road in the north. The park occupies approximately



**FIGURE 1**  
Location of the Highveld National Park in North-West Province, South Africa, with the insert placing the HNP in the context of the rest of South Africa

10 200 hectares, which consists of a core area of Potchefstroom local authority land (5500 ha), Modderfontein (Defence Force Land, 2 700 ha), Nooitverwacht (privately owned land, 1500 ha) and the Agricultural College Farm (500 ha).

### Geology, soil and land types

The relatively heterogeneous geology is represented by the Witwatersrand and Ventersdorp supergroups and the Transvaal Sequence, with isolated occurrences of old Archaic granites and Karoo Sequence sediments (Bezuidenhout 1993). Soils in the study area are heterogeneous and vary from sandy to clayey, due to great variation in parent rock material (Land Type Survey Staff 1984). Four land types – Ba, Bc, Fa and Fb – occur in the HNP (Daemane 2007). A land type denotes an area that can be shown at a scale of 1:250 000, displaying marked degree of uniformity with respect to terrain form, soil pattern and climate (Land Type Survey Staff 1984).

### Climate

The rainfall is erratic and varies from an average of 600 mm per annum, to exceptional occurrences of more than 900 mm per annum (Institute for Soil, Climate and Water 2003). The area is characterised by large seasonal and daily variations in temperature, from very hot in summer (daily average temperatures may exceed 32 °C in January), to mild to cold in winter, with average minimum monthly temperatures of up to 12 °C (ISCW 2003).

### Vegetation

The North-West Province encloses two major biomes: the Grassland and the Savanna Biome. The Grassland Biome in this study was represented by the Vaal Reefs Dolomite Sinkhole Woodland (Gh 12), the Rand Highveld Grassland (Gm 11), the Klerksdorp Thornveld (Gh 13), and the Carletonville Dolomite

Grassland (Gh 15) of the Dry Highveld Grassland Bioregion Unit (Mucina & Rutherford 2006). The Savanna Biome, represented by the Central Bushveld Bioregion Unit, consists of Andesite Mountain Bushveld (SVcb 11). Low & Rebelo (1996) classified this area under the Rocky Highveld Grassland and Dry Sandy Highveld Grassland. The Bankenveld vegetation type (Veld Type 61), which Acocks (1953) described as ‘false grassveld’, is dominant in the HNP; the climax of this veld type is open *Acacia caffra* woodland.

## METHOD


### Stratification of sample sites

Vegetation sampling was undertaken to represent the communities of the HNP and consisted of 88 relevés of 900 m<sup>2</sup>, each randomly placed within the stratified units. Stratification was done on 1:50 000 scale aerial orthophotographs, on the basis of relatively homogeneous physiographic and physiognomic units. Following Bezuidenhout (1993), further stratification was carried out by recognising the terrain types associated with topographical positions, such as the crest, midslope, footslope, valley bottomlands and floodplains.

### Sampling procedure

Plant species were identified in each relevé and the cover abundance of each species was visually estimated using the Braun-Blanquet Cover Abundance Scale (Mueller-Dombois & Ellenberg 1974). Unknown specimens were identified in the Kimberley South African National Parks Herbarium. Germishuizen & Meyer (2003) was used as reference to confirm the names of taxa. In addition, the average height and canopy cover of tree, shrub and herbaceous strata were estimated. Features of the habitat were recorded for each relevé, including geology, soil forms (MacVicar *et al.* 1991), and rockiness of the soil surface





**FIGURE 2**  
Vegetation map of the plant communities of the Highveld National Park, North-West Province, South Africa

**Data processing**

The TWINSpan classification algorithm (Hill 1979), which is regarded as a very successful approach by many phytosociologists (Mucina & Van der Maarel 1989; Bredenkamp & Bezuidenhout 1995; Cilliers 1998), was used as a first analysis for the floristic data. The TWINSpan classification was then refined further, by application of Braun-Blanquet procedures by means of the BBPC-program (Bezuidenhout, Biggs & Bredenkamp 1996). Detrended correspondence analysis (DCA; Hill 1979) was applied to the data set to confirm the phytosociological association and to assess floristic relations between communities. The stepwise ordination approach followed by Mucina & Van Tongeren (1989), was adopted to cope with the heterogeneity of the data set. Descriptive terms such as ‘differential’ and ‘dominant species’ were used following Kent & Coker (1992) and Jennings *et al.* (2009). ‘Differential species’ refers to species of medium to low constancy, which tends to occur together in a series of quadrats and can thus be used to characterise groups. ‘Dominant species’ refers to species with the highest percentage of cover, usually in the uppermost dominant layer.

**RESULTS**

**Vegetation classification**

The analysis of vegetation by phytosociological means, resulted in the identification of nine different plant communities (Figure

2), brief descriptions of which can be found in Table 1. The species groups within each plant community can be found in the phytosociological table (see Online Appendix). The presence of the municipal dumping site within the park is shown in the map (Figure 2) and its implications are highlighted in the discussion section. However, the area indicated as ‘Spitskop’ on the map is highly degraded; therefore, the plant communities found in this area do not form part of this discussion.

**Detailed description of plant communities for HNP**

**1. *Mystroxyloa aethiopicum* – *Pavetta zeyheri* shrubland.** This community is mainly limited to the rocky quartzite plateau and ridges of the study area. Rocks cover up to 73% of the soil surface. Glenrosa is the dominant soil form in this community. The differential species of this community include indigenous species such as *Pavetta zeyheri*, *Mystroxyloa aethiopicum*, *Grewia occidentalis*, *Vangueria parvifolia*, *Pappea capensis*, *Olea europaea* subsp. *africana* and *Panicum maximum* (Species Group A). Dominant species include *Themeda triandra*, *Elionurus muticus* (Species Group U), *Vangueria infausta* subsp. *infausta* (Species Group C), and *Acacia caffra* (Species Group N). The average height of the tree stratum is 5.3 m, with a canopy cover of 19%. The average height of the herbaceous stratum is 0.9 m, with a canopy cover of 41%. On average, 33 species were recorded per sample plot, of which 1.2% was exotic species. Exotic plants included species such as *Gomphrena celosioides* and *Opuntia ficus-indica* (Species Group V).

**TABLE 1**  
A summary of the plant communities described for the Highveld National Park

Plant community	Brief description of location
1. <i>Myroxylon aethiopicum</i> – <i>Pavetta zeyheri</i> shrubland	Mainly limited to the plateaus of the andesitic lava and rocky quartzite hills
2. <i>Vangueria infausta</i> subsp. <i>infausta</i> – <i>Loudetia simplex</i> shrubland	Associated with the midslopes of the rocky quartzite ridges
3. <i>Diheteropogon amplexens</i> – <i>Trachypogon spicatus</i> grassland	Occurs on stony plains and midslopes of quartzite hills and andesitic lava
4. <i>Schizachyrium sanguineum</i> – <i>Cymbopogon excavatus</i> grassland	Associated with the midslopes of the chert outcrops
5. <i>Rhus pyroides</i> – <i>Acacia erioloba</i> woodland	Limited to the valley bottomlands of the chert and dolomitic areas
6. <i>Acacia caffra</i> – <i>Setaria sphacelata</i> var. <i>sphacelata</i> woodland	Associated with the midslopes of the andesitic lava and quartzite hills
7. <i>Acacia karroo</i> – <i>Ziziphus zeyheriana</i> woodland	Occurs on the footslopes, valley bottomlands and along the drainage lines
8. <i>Cymbopogon pospischilii</i> – <i>Themeda triandra</i> grassland	Associated with the midslopes of the quartzite and chert outcrops
9. <i>Setaria incrassata</i> – <i>Hyparrhenia hirta</i> grassland	Occurs in the quartzite and colluvial bottomlands or floodplains in or along drainage lines

A closely related shrubland community, *Pavetta zeyheri* – *Vangueria infausta* of the Ba land type of the former western Transvaal, was described by Bezuidenhout, Bredenkamp and Theron (1994c) and corresponds with the Dry Sandy Highveld Grassland. The shrubland communities form a distinctive structural vegetation type within the matrix of the Dry Sandy Highveld Grassland. Only 0.28% of the Dry Sandy Highveld Grassland is currently being conserved in South Africa, while approximately 65% of this vegetation has been transformed (Bredenkamp & Van Rooyen 1996b).

**2. *Vangueria infausta* subsp. *infausta* – *Loudetia simplex* shrubland.** This community is associated with the midslopes of the rocky quartzite outcrops of the study area. The Mispah and Glenrosa soil forms are dominant in this habitat. Rocks cover up to 63% of the soil surface. The differential species of this community is Species Group B, which includes the grass *Loudetia simplex*, the shrub *Mundulea sericea* and a forb *Helichrysum kraussii*. The dominant shrub is *Vangueria infausta* subsp. *infausta* (Species Group C). The dominant grasses are *Trachypogon spicatus* and *Andropogon schirensis* (Species Group E), *Schizachyrium sanguineum*, *Brachiaria serrata*, *Diheteropogon amplexens*, *Eragrostis racemosa*, *Melinis nerviglumis*, and *Aristida meridionalis* (Species Group F), and *Setaria sphacelata* var. *sphacelata* and *Eragrostis curvula* (Species Group T). The average height of the tree stratum is 4.6 m and the average canopy cover is 6.8%, with *Acacia caffra* (Species Group N) forming the major tree component. The average canopy cover of the herbaceous stratum is 53.6%, with an average height of 0.8 m. On average, 34 species were recorded per sample plot, of which 1.6% was exotic species. Exotic plants included species such as *Tagetes minuta* (Species Group L), *Verbena officinalis* (Species Group S), and *Conyza bonariensis*, *Achyranthes aspera* and *Nicotiana longiflora* (Species Group V).

Cilliers, Schoeman & Bredenkamp (1999) described a closely related shrubland community (*Vangueria infausta* subsp. *infausta* – *Rhus pyroides*) in the semi-urban area of Potchefstroom. According to their findings, the lack of the *Acacia caffra* and the dominance of *Rhus pyroides* in the semi-urban area of Potchefstroom is possibly a result of the encroachment of *Rhus pyroides* from the adjacent dolomitic and chert grasslands of the Fa land type. A similar community, the *Vangueria infausta* subsp. *infausta* – *Acacia caffra* woodland was described for the Bc land type of the former western Transvaal Grassland (Bezuidenhout & Bredenkamp 1991). According to Bezuidenhout & Bredenkamp (1991), the Bc land type can be divided into two broad physiognomic classes, namely woodland and grassland. The woodland competes with the grassland for more or less the same habitat. The grassland is often overgrazed and burned and the dominance of *Mundulea sericea* is indicative of these disturbances. These shrubland communities, typical of the Rocky Highveld Grassland (Bredenkamp & Van Rooyen 1996a), are well worth conserving.

**3. *Diheteropogon amplexens* – *Trachypogon spicatus* grassland.** This grassland occurs mostly on the rocky plains and midslopes.

The geology in this community is characterised by quartzite and andesitic lava. Rocks cover 48% of the soil surface. The soil forms comprise the Mispah and Glenrosa. Differential species of this community, indicated by Species Group D, include the grass – *Brachiaria nigropedata*, the forbs – *Helichrysum nudifolium* var. *nudifolium* and *Chascanum adenostachyum*, and the succulent – *Aloe zebrina*. The dominant grass species in this community are *Diheteropogon amplexens* (Species Group F) and *Trachypogon spicatus* (Species Group E). The average height of the herbaceous stratum is 0.9 m and the canopy cover is 65%. The average height of the tree stratum is 4.6 m, with a canopy cover of 3.8%; *Acacia caffra* (Species Group N) is the main component of this stratum. The average height of the shrub stratum is 1.5 m, with a canopy cover of 9%. On average, 33 species were recorded per sample plot, with the only exotic species (0.2%) found in this community being *Conyza bonariensis* (Species Group V).

A similar plant community, the *Trachypogon spicatus* – *Diheteropogon amplexens* grassland from the dolomitic region in the Potchefstroom–Ventersdorp–Randfontein area, was previously described by Bezuidenhout & Bredenkamp (1990). Other closely related communities described by Bezuidenhout, Bredenkamp and Theron (1994b) include the *Diheteropogon amplexens* – *Schizachyrium sanguineum* grassland from the Fb land type. Bezuidenhout *et al.* (1994b) also found that the *Diheteropogon amplexens* – *Trachypogon spicatus* grassland occurs in the Fb land type and, although it is still relatively undisturbed, it is highly threatened by mining and industrial activities elsewhere and is therefore worth conserving. Cilliers *et al.* (1999) described the *Schizachyrium sanguineum* – *Diheteropogon amplexens* grassland in the natural and semi-natural areas in the municipal area of Potchefstroom, noting that the variation between these previously described communities could be contributed to factors such as altitude, terrain type, soil type, soil depth and rockiness of soil surface.

**4. *Schizachyrium sanguineum* – *Cymbopogon excavatus* grassland.** This community is associated with the midslopes of the chert outcrops, where Mispah is the dominant soil form. Rocks cover 37% of the soil surface. No differential species were identified in this community, but it is characterised by the absence of species within Species Groups A and C. The dominant species are *Schizachyrium sanguineum*, *Cymbopogon excavatus*, *Brachiaria serrata*, *Diheteropogon amplexens*, and *Dicoma anomala* (Species Group F), *Heteropogon contortus* (Species Group Q), *Eragrostis curvula* (Species Group T), and *Pseudognaphalium undulatum* (Species Group S). The average height of the herbaceous stratum is 0.7 m and it has a canopy cover of 66.3%. The average height of the tree stratum is 6.8 m, with only a poor canopy cover of 1.5%. *Celtis africana* and *Acacia erioloba* (Species Group G) are the only tree species in this community and are very poorly represented. On average, 54 species were recorded per sample plot, of which 0.7% was exotic species. Exotic plants included species such as *Conyza bonariensis* and *Achyranthes aspera* (Species Group V).

Bezuidenhout & Bredenkamp (1990) described a closely related community, *Aristida diffusa* – *Cymbopogon excavatus* grassland, as





also occurring on the stony chert areas of the dolomitic region in the Potchefstroom–Ventersdorp–Randfontein. Another related community, *Cymbopogon excavatus* – *Diheteropogon amplexans* Grassland, found in the former western Transvaal dolomite and chert grassland, was described by Bezuidenhout, Bredenkamp and Theron (1994a). He associated this grassland with the high-lying areas of the Fa land type, which has a high conservation status due to its floristic and habitat diversity.

**5. *Rhus pyroides* – *Acacia erioloba* woodland.** This woodland is limited to the valley bottomlands of the chert and dolomitic areas. Mispah is the dominant soil form in this community, with rocks forming 30% of the soil surface. Sinkholes, underlain by dolomite, are present in this habitat and mostly filled by deep sandy soil (> 1.2 m). Differential species of this community, indicated by Species Group G, include *Eragrostis superba*, *Sporobolus africana*, *Eragrostis trichophora*, *Celtis africana*, *Acacia erioloba*, *Diospyros lycioides* subsp. *guerkei*, *Ipomoea oblongata*, *Sida dregéi*, and *Striga asiatica*. Dominant species in this plant community are *Acacia erioloba* (Species Group G), *Triraphis andropogonoides* (Species Group H), *Ehretia rigida* (Species Group L), *Rhus pyroides* (Species Group M), and *Cymbopogon pospischilii* (Species Group Q). The average height of the tree stratum is 5.7 m, with a canopy cover of 4.8%. The trees are restricted to deep sandy soils in the sinkholes. The average height of the shrub stratum is 2.8 m, with a canopy cover of 15.6 m. The average height of the herbaceous stratum is 0.8 m, with a high canopy cover of 67%. The succulent *Lithops lesliei* is also found in this community (Species Group V). On average, 40 species were recorded per sample plot, of which 4.5% was exotic species. Exotic plants included *Tagetes minuta* (Species Group L) and *Conyza bonariensis*, *Achyranthes aspera*, *Bidens bipinnata*, and *Solanum nigrum* (Species Group V).

A closely related woodland community, *Rhus lancea* – *Acacia erioloba* in the former western Transvaal (Bezuidenhout *et al.* 1994b), shares a similar floristic composition with the *Rhus pyroides* – *Acacia erioloba* woodland, except for the dominance of *Rhus pyroides* in the current study. The *Rhus* – *Acacia* woodland is aesthetically associated with the most scenic landscapes in the western Grassland Biome and certainly deserves high conservation priority (Bezuidenhout 1993). This community also falls within the Vaal Reefs Dolomite Sinkhole woodland (Gh 12), which occurs naturally in clumps around sinkholes, especially in places of dolomite outcrops (Mucina & Rutherford 2006). The HNP is supposed to conserve a considerable area of this vegetation unit (Mucina & Rutherford 2006). Gathering of wood for fuel by rural poor communities is the most significant threat to this plant community.

**6. *Acacia caffra* – *Setaria sphaelata* var. *sphaelata* woodland.** The *Setaria sphaelata* var. *sphaelata* – *Acacia caffra* woodland is mainly associated with the andesitic lava and quartzite midslopes of the rocky outcrops in the study area. Rocks cover 27% of the soil surface. Glenrosa is the dominant soil form, with Mispah and Hutton to a lesser extent. The differential species in this community are found in Species Group I and include the grasses – *Digitaria eriantha*, *Eragrostis rigidior* and *Aristida transvaalensis*, the forbs – *Crabbea angustifolia*, *Blepharis integrifolia*, *Indigofera heterotricha* and *Chaetacanthus costatu*, the tree *Euclea undulate*, and the shrub *Rhus leptodictya*. Dominant species in this community include *Setaria sphaelata* var. *sphaelata* (Species Group T), *Acacia caffra* (Species Group N), *Pseudognaphalium undulatum*, *Aloe greatheadii* var. *Davyana*, and *Clematis brachiata* (Species Group N), *Ziziphus zeyheriana* (Species Group Q), and *Acacia karroo* (Species Group S). The average height of the tree stratum is 5 m, with a canopy cover of 11.2%. The average height of the shrub stratum is 1.9 m, with a canopy cover of 17.9%. The average height of the herbaceous stratum is 0.9 m, with a canopy cover of 61.7%. On average, 40 species were recorded per sample plot, of which 6.6% was exotic species. Exotic plants included species such as *Tagetes minuta* (Species Group L), *Verbena officinalis* (Species Group S), *Paspalum dilatatum*, *Conyza*

*bonariensis*, *Achyranthes aspera*, *Bidens bipinnata*, *Gomphrena celosioides*, *Schkuhria pinnata*, *Verbena bonariensis*, *Gulleminea densa*, *Plantago lanceolata*, *Solanum sisymbriifolium*, *Chenopodium album*, *Opuntia ficus-indica*, *Oxalis corniculata*, *Zinnia peruviana*, *Oenothera tetraptera*, and *Cirsium vulgare* (Species Group V).

Bezuidenhout *et al.* (1994b) described a closely related community, *Rhus leptodictya* – *Acacia caffra* woodland of the Fb land type in the former western Transvaal Grassland. Another related community (*Vangueria infausta* – *Acacia caffra* woodland) of the Bc land type in the former western Transvaal Grassland, was described by Bezuidenhout & Bredenkamp (1991). These communities correspond with the Andesite Mountain Bushveld (SVcb 11), described by Mucina & Rutherford (2006), which is dominated by species such as *Acacia karroo*, *Acacia caffra* and *Ziziphus zeyheriana*. Intensive mining of the gold-bearing quartzitic rocks of the Witwatersrand supergroup has caused destruction and degradation to much of the vegetation within the Fb land type (Bezuidenhout 1993). Uncontrolled fires, gathering of trees for fuel wood and accompanying disturbances, such as trampling and soil compaction, are also significant threats to these communities (Cilliers *et al.* 1999).

**7. *Acacia karroo* – *Ziziphus zeyheriana* woodland.** The *Acacia karroo* – *Ziziphus zeyheriana* Woodland occurs on footslopes, valley bottomland areas and along the drainage lines and, occasionally, on the midslopes in degraded areas. Rocks cover 3% of the soil surface, with Glenrosa and Hutton as the dominant soil forms in this community (although Mispah and Shortland occur in few relevés). The geology is mainly quartzitic, with andesitic lava in few relevés. The differential species in this community include *Delosperma herbeum*, *Nidorella anomala*, *Sida spinosa*, *Pavonia burchellii*, *Pollichia campestris*, *Pentzia globosa* and *Eragrostis lehmanniana* (Species Group J). Dominant species are *Ziziphus zeyheriana* (Species Group Q), *Acacia karroo* (Species Group S), *Asparagus suaveolens* (Species Group K), *Ehretia rigida* (Species Group L), *Aloe greatheadii* var. *davyana*, *Clematis brachiata*, and *Acacia caffra* (Species Group N), *Cymbopogon pospischilii* and *Teucrium trifidum* (Species Group Q), *Cynodon dactylon* and *Asparagus larinus* (Species Group S), and *Setaria sphaelata* var. *sphaelata* (Species Group T). The average height of the tree stratum is 4.5 m and the canopy cover is 17.2%. The average height of the shrub stratum is 1.8 m, with a canopy cover of 21.9%. The average height of the herbaceous stratum is 0.9 m, with a canopy cover of 6.5%. On average, 41 species were recorded per sample plot, with 7.2% exotic species. Exotic plants included species such as *Tagetes minuta* (Species Group L), *Verbena officinalis* (Species Group S), *Conyza bonariensis*, *Achyranthes aspera*, *Bidens bipinnata*, *Gomphrena celosioides*, *Lepidium bonariensis*, *Schkuhria pinnata*, *Solanum sisymbriifolium*, *Opuntia ficus – indica*, *Oxalis corniculata*, and *Zinnia peruviana* (Species Group V).

Bredenkamp, Joubert and Bezuidenhout (1989) described a similar community (*Acacia karroo* woodland) in the Potchefstroom–Fochville–Parys area, with *Ziziphus zeyheriana* abundantly occurring in severely degraded areas. They also stated that this community is an encroachment of the *Acacia karroo* woodland into grassland communities. The conservation value of this vegetation type, based on the variety in habitat and floristic diversity, was emphasised by Bezuidenhout *et al.* (1994b) in the syntaxonomy of the vegetation of the Fb land type in the former western Transvaal Grassland.

**8. *Cymbopogon pospischilii* – *Themeda triandra* grassland.** This grassland is associated with the rocky quartzite midslopes of the study area. Rocks cover 3% of the soil surface, with Glenrosa, Hutton and Shortland as the dominant soil forms. The differential species in this community are in Species Group O and include species such as *Hibiscus pusillus*, *Convolvulus sagittatus*, *Eragrostis chloromelas*, *Menodora africana*, *Gladiolus splendens*, *Helichrysum rugulosum*, *Gymnosporia buxifolia*, and *Anthospermum hispidulum*. Dominant species in this community are *Cymbopogon*

*pospischilii* and *Ziziphus zeyheriana* (Species Group Q), *Themeda triandra* (Species Group U) *Hermannia depressa* (Species Group P), *Berkheya radula*, *Hyparrhenia hirta*, *Setaria incrassata*, and *Helichrysum nudifolium* var. *nudifolium* (Species Group R), and *Acacia karroo* and *Conyza podocephala* (Species Group S). The average height of the tree stratum is 5.4 m, with a canopy cover of 9.8%. The average height of the shrub stratum is 1.2 m, with a canopy cover of 8.2%. The average height of the herbaceous stratum is 0.9 m, with a canopy cover of 77%. On average, 31 species were recorded per sample plot, of which 6.3% was exotic species. Exotic plants included species such as *Tagetes minuta* (Species Group L) *Verbena officinalis* (Species Group S) *Conyza bonariensis*, *Achyranthes aspera*, *Avena fatua*, *Bidens bipinnata*, *Gomphrena celosoides*, *Physalis angulata*, *Oenothera rosea*, *Oenothera tetraptera*, and *Cirsium vulgare* (Species Group V).

A closely related plant community (*Themeda triandra* – *Elyonurus muticus* grassland) with an absolute dominance of *Themeda triandra*, was described by Bredenkamp, Joubert and Bezuidenhout (1989) in a reconnaissance survey of the vegetation of the plains in the Potchefstroom–Fochville–Parys areas. In the Bc land type, the *Cymbopogon pospischilii* – *Themeda triandra* grassland shares floristic characteristics with the *Themeda triandra*, a variant described by Bezuidenhout & Bredenkamp (1991), who found vegetation associated with these plant communities to be in a state of degradation, due to continued overgrazing by domestic stock.


**9. *Setaria incrassata* – *Hyparrhenia hirta* grassland.** The *Setaria incrassata* – *Hyparrhenia hirta* grassland is restricted to the valley bottomlands and floodplains, and in or along drainage lines within the study area. The geology comprises mostly colluvial deposits from the upper slopes. Rocks cover 1% of the soil surface. This habitat is fairly unstable due to seasonal

flooding and drying. Hutton and Swartland are the dominant soil forms found in this habitat. No differential species were identified in this community. *Setaria incrassata* and *Hyparrhenia hirta* (Species Group R) are the dominant species. The average height of the herbaceous stratum is 0.9 m, with a high canopy cover of 69%. The average height of the shrub stratum is 0.8 m, with a poor canopy cover of 1.3%. The average height of the tree stratum is 3 m and a canopy cover of 6%. On average, 24 species were recorded per sample plot, of which 12% was exotic species. Exotic plants included *Verbena officinalis* (Species Group S), and *Paspalum dilatatum*, *Conyza bonariensis*, *Ciclospermum leptophyllum*, *Persicaria lapathifolia*, *Oenothera rosea* and *Oenothera tetraptera* (Species Group V).

Several researchers have done studies on similar wetland grassland communities. Bredenkamp *et al.* (1994) described a *Hyparrhenia hirta* grassland in the Boskop Dam Nature reserve, Bezuidenhout *et al.* (1994b) described a *Hyparrhenia hirta* – *Eragrostis plana* grassland in the Fb land type in the former western Transvaal, and Cilliers *et al.* (1998) gave an account of a *Hyparrhenia hirta* grassland in the Potchefstroom municipality. The similarity found in all these communities was that their habitats were fairly unstable due to seasonal flooding and drying which, together with the frequent overgrazing of these sites, caused an advanced state of degradation of the vegetation.

**Ordination**

DCA was applied to the data set to confirm the existence of the different plant communities and to determine whether the gradients in the plant communities along different ordination axes could be explained by gradients in specific environmental characteristics. The gradient along ordination axis 1 (E = 0.65) could be explained by surface rockiness (Figure 3). Plant communities to the left of the scatter diagram (plant



**FIGURE 3** Ordination of 87 relevés in Highveld National Park showing the relationship between surface rockiness and altitude



communities 1 and 2) are characterised by a largely rocky soil surface (> 50%), while those on the right (plant communities 7, 8 and 9) are characterised by fewer rocks on the soil surface (0% – 25%). Plant communities at the centre of the scatter diagram (plant communities 3, 4, 5 and 6) are characterised by a moderately rocky surface (25% – 50%). The gradient along ordination axis 2 ( $E = 0.45$ ) could be explained by altitude. Plant communities at the top of the scatter diagram occurred mainly on the rocky quartzite plateau and crests in the HNP. Plant communities found at the bottom of the scatter diagram are associated with relatively low altitudes and occur mainly in the valley bottomlands of the cherts and dolomitic areas.

## DISCUSSION

Three structural vegetation units, namely woodland, shrubland and grassland, were identified in the HNP. The DCA-ordination also confirmed the relationship between the plant communities and the environmental gradients. Many of the plant communities previously described in the grassland of the former western Transvaal also show floristic and habitat affinities with the plant communities described in the present study. The *Acacia karroo* woodland, as well as several grassland communities, occurs on the plains, while *Acacia caffra* woodland occurs on the rocky outcrops and hills (Bezuidenhout & Bredenkamp 1991; Bezuidenhout *et al.* 1994a; Bezuidenhout *et al.* 1994b). The plant communities described in the present study also fall within the poorly conserved Rocky Highveld and Dry Sandy Highveld Grasslands previously described by Bredekamp & Van Rooyen (1996a; 1996b). These vegetation types are highly threatened by urbanisation, mining and agriculture. The HNP aims to conserve a considerable area of the Vaal Reefs Dolomite Sinkhole Woodland (Gh 12), because, according to Mucina and Rutherford (2006), this is one of the most scenic landscapes in the western Grassland Biome. Mucina and Rutherford (2006), also point out that almost a quarter of this vegetation unit has already been transformed by mining (the region contains possibly the highest concentration of mines in South Africa), cultivation, urban sprawl, and road construction.

Five alien species belonging to Category 1 plants were identified in the HNP. Category 1 plants are described as prohibited species that should not be tolerated because they are able to invade undisturbed environments and transform or degrade natural plant communities (Henderson 2001). These five species are *Opuntia ficus-indica* (Species Group S), and *Achyranthes aspera*, *Cirsium vulgare*, *Solanum elaeagnifolium*, and *Solanum sisymbriifolium* (Species Group V). Focus should be given to plant communities 6 and 7, as they both have the highest number of category 1 species, with 6.6% and 7.2 % per sample plot, respectively. Even though plant community 9 has the highest number of alien species per sample plot (12%), no Category 1 species were found in this community. Control of these species should aim towards their long-term reduction in density and abundance. It is therefore crucial to prevent the further spread of alien plants into un-invaded areas by isolating the dense patches within the landscape through appropriate control measures and monitoring.

The unique scenario of the existence of the municipal dumping site within the park is a cause for concern and it cannot be emphasised enough that proper management and control of the dump site is crucial. Landfill exfiltration and water leakage can adversely affect the quality of the adjoining soils and consequently affect the flora and fauna in the adjoining areas. Air pollution and wind-carried suspensions can also have direct effects in the park. Although the current operations ensure that storm water drains away from the waste to prevent contamination of any runoff, the closure and rehabilitation of the dump is recommended. The aesthetics of the park are compromised by the dump, its associated infrastructure and heavy traffic.

The HNP is also threatened by increasing urbanisation, which is why it is crucial to recognise the threats to biodiversity and to understand the factors that will influence the management of the

park. For example, settlements along the park's border pose a threat to the park on a number of levels, including giving rise to noise, air and land pollution, the loss of wilderness ambience, as well as representing a sustained pressure on natural resources. South African cities, including Potchefstroom, have witnessed an increase in the number of informal settlements on their urban fringe. There has also been an increase in squatting and informal settlements in rural areas, where fewer job opportunities exist and higher poverty levels are prevalent (Cilliers *et al.* 2004). Incidences such as fence cutting, harvesting of trees for firewood, uncontrolled fires and overgrazing, are just a few of the challenges that face the HNP. Therefore research on the HNP should focus not only on those problems that can be measured ecologically, but also those that allow for sustainable resource use. This highlights the importance of conservation to the economy in general, and to people's lives, in particular. The rural and urban poor are those most affected because they depend directly on the natural environment for their livelihood. Considerable areas covered by plant communities have not only been transformed by urban sprawl, but existing plant communities on the urban fringe of Potchefstroom are also highly fragmented (Cilliers *et al.* 2004).

## CONCLUSION

The fundamental role of protected areas has been to separate elements of biodiversity from processes that threaten their existence, by establishing them in remote areas (Margules & Pressey 2000). Recently, however, a more systematic approach to locating and designing protected areas has been evolving – one in which the influences of communities are taken into consideration. This approach needs to be implemented if a large proportion of today's biodiversity is to endure in the future. The park has not yet been proclaimed, but is currently under the management of North-West Parks and Tourism Board. It will therefore require a management plan that will provide an overarching spatial planning framework. Such a framework should address issues regarding urban development close to the park, as well as its future expansion. The ecological interpretation derived from this study can be used as a tool for environmental planning, management and conservation of this grassland area.

## ACKNOWLEDGEMENTS

The following individuals and institutions are sincerely thanked: South African National Parks for financing this study, North-West University for supporting this research, Dr. Stephen Holness and Ms. Phozisa Mamfengu for assisting with maps, and Mr. Abbey Legari for assisting with field work and the Highveld National Park' Founding Partners for allowing us to undertake this study.

Editor: Llewellyn C. Foxcroft

## REFERENCES

- Acocks, J.P.H., 1953, 'Veld types of South Africa', *Memoirs of the Botanical Survey of South Africa* 28, 1–192.
- Archer, F., 1999, *Participatory decision-making at Richtersveld Contractual Park: Myth or reality?*, draft report to the Evaluating Eden Project, International Institute for Environment and Development, London.
- Bezuidenhout, H., & Bredenkamp, G.J., 1991, 'The vegetation of the Bc land type in the western Transvaal Grassland, South Africa', *Phytocoenologia* 19(4), 497–518.
- Bezuidenhout, H., 1993, 'Syntaxonomy and synecology of western Transvaal grassland', PhD thesis, Department of Botany, University of Pretoria.
- Bezuidenhout, H. & Bredenkamp, G.J., 1990, 'A reconnaissance survey of the vegetation of the dolomitic region in the Potchefstroom–Ventersdorp–Randfontein area, South Africa', *Phytocoenologia* 18 (2/3), 387–403.
- Bezuidenhout, H., Bredenkamp, G.J. & Theron, G.K., 1994a, 'A classification of the vegetation of the western Transvaal dolomite and chert grassland, South Africa', *South African Journal of Botany* 60(3), 152–161.



- Bezuidenhout, H., Bredenkamp, G.J. & Theron, G.K., 1994b, 'The syntaxonomy of the vegetation of the Fb land type in the western Transvaal Grassland, South Africa', *South African Journal of Botany* 60(3), 72–81.
- Bezuidenhout, H., Bredenkamp, G.J. & Theron, G.K., 1994c, 'The vegetation syntaxa of the Ba land type in the western Transvaal Grassland, South Africa', *South African Journal of Botany* 60(4), 214–224.
- Bezuidenhout, H., Biggs H.C. & Bredenkamp, G.J., 1996, 'A process supported by the utility BBPC for analysis of Braun-Blanquet data on personal computer', *Koedoe* 39, 107–112.
- Bredenkamp, G.J. & Bezuidenhout, H., 1989, 'A phytosociology of the Faan Meintjies Nature Reserve in the western Transvaal grassland, South Africa', *Die Suid Afrikaanse Tydskrif vir Natuurwetenskap en Tegnologie* 56(1), 54–64.
- Bredenkamp, G.J., Joubert, A.F. & Bezuidenhout, H., 1989, 'A reconnaissance survey of the vegetation of the plains in the Potchefstroom–Fochville–Parys areas', *South African Journal of Botany* 55(2), 199–206.
- Bredenkamp, G.J., Bezuidenhout, H., Joubert, H. & Naude, C., 1994, 'The vegetation of the Boskop Dam nature Reserve, Potchefstroom', *Koedoe* 37(1), 19–33.
- Bredenkamp, G. J. & Bezuidenhout, H., 1995, 'A proposed procedure for the analysis of large data sets in the classification of South African Grasslands', *Koedoe* 38(1), 33–39.
- Bredenkamp, G.J. & Van Rooyen, N., 1996a, 'Rocky Highveld Grassland', in A.B. Low & A.G. Rebelo (eds.), *Vegetation of South Africa, Lesotho and Swaziland*, p. 39, Department of Environmental Affairs and Tourism, Pretoria.
- Bredenkamp, G.J. & Van Rooyen, N., 1996b, 'Dry Sandy Highveld Grassland', in A.B. Low & A.G. Rebelo (eds.), *Vegetation of South Africa, Lesotho and Swaziland*, p. 41, Department of Environmental Affairs and Tourism, Pretoria.
- Cilliers, S.S., 1998, 'A phytosociological study of urban open spaces in Potchefstroom, North-West Province, South Africa', PhD thesis, School of Environmental Sciences and Development, Potchefstroom University for Christian Higher Education.
- Cilliers, S.S., Schoeman, L.L. & Bredenkamp, G.J., 1998, 'Wetland plant communities in the Potchefstroom municipal area, North-West Province, South Africa', *Bothalia* 28(2), 213–229.
- Cilliers, S.S. & Bredenkamp, G.J., 1998, 'Vegetation analysis of railway reserves in the Potchefstroom municipal area, North-West Province, South Africa', *South African Journal of Botany* 64(5), 271–280.
- Cilliers, S.S., Van Wyk, E. & Bredenkamp, G.J., 1999, 'Urban nature conservation: Vegetation of natural areas in the Potchefstroom municipal area, North-West Province, South Africa', *Koedoe* 42(1), 1–30.
- Cilliers, S.S., Müller, N. & Drewes, E., 2004, 'Overview on urban nature conservation: situation in the western-grassland biome of South Africa', *Urban Forestry & Urban Greening* 3, 49–62.
- Daemane, M.E., 2007, 'An ecological study of the plant communities and degraded areas of the Highveld National Park, North-West Province, South Africa', MSc thesis, School of Environmental Sciences and Development, North-West University.
- Edwards, D., 1983, 'A broad-scale structural classification of vegetation for practical purposes', *Bothalia* 14(3/4), 705–712.
- Fabricius, C., Koch, E. & Magome, H., 2001, 'Towards strengthening collaborative ecosystem management: Lessons from environmental conflict and political changes in southern Africa', *Journal of the Royal Society of New Zealand* 31(4), 831–844.
- Fairbanks, D.H.K., Thomson, M.W., Vink, D.E., Newby, T.S., Van den Berg, H.M. & Everard, D.A., 2000, 'The South African land-cover characteristics database: A synopsis of the landscape', *South African Journal of Science* 96, 69–82.
- Germishuizen, G. & Meyer, N.L. (eds.), 2003, 'Plants of Southern Africa: An annotated checklist', *Strelitzia* 14, 1–1231.
- Hill, M.O., 1979, *TWINSPAN - A FORTRAN program for the arranging multivariate data in an ordered two-way table by classification of individuals and attributes*, Cornell University, New York.
- Hoffman, T. & Todd, S., 2000, 'A national review of land degradation in South Africa: The influence of biophysical and socio-economic factors', *Journal of Southern African Studies* 26(4), 743–758.
- Hoffman, M.T. & Meadows, M.E., 2002, 'The nature, extent and causes of land degradation in South Africa: Legacy of the past, lessons for the future?', *Area* 34(4), 428–437.
- Institute for Soil, Climate and Water, 2003, 'Comprehensive Report: Brakspruit, Potchefstroom', Agricultural Research Council, Agromet, Pretoria.
- Jennings, M.D., Faber-Langendoen, D., Loucks, O.L., Peet, R.K. & Roberts, D., 2009, 'Standards for associations and alliances of the U.S. National Vegetation Classification', *Ecological Monographs* 79, 173–199.
- Kent, M. & Coker, P., 1992, *Vegetation description and analysis: A practical approach*, John Wiley & Sons Ltd., Chichester.
- Land Type Survey Staff, 1984, 'Land types of the maps 2626 West Rand and 2726 Kroonstad', *Memoirs of Agricultural Natural Resources of South Africa* 4, 1–441.
- Low, A.B. & Rebelo, A.G., 1996, *Vegetation of South Africa, Lesotho and Swaziland*, Department of Environmental Affairs and Tourism, Pretoria.
- Macvicar, C.N., Loxton, R.F., Lambrechts, J.J.N. et al., 1991, *Soil Classification a taxonomic system for South Africa*, Department of Agricultural Development, Soil and Irrigation Research Institute, Pretoria. *Memoirs on the Agricultural Natural Resources of South Africa* 15.
- Margules, C.R. & Pressey R.L., 2000, 'Systematic conservation planning' *Nature* 405, 243–253.
- Mucina, L. & Van Tongeren, O.F.R., 1989, 'A coenocline of the high-ranked syntaxa of ruderal vegetation', *Vegetation* 81, 117–125.
- Mucina, L. & Rutherford, M.C. (eds.), 2006, 'The vegetation of South Africa, Lesotho and Swaziland', *Strelitzia* 19, 1–807.
- Mucina, L. & Van der Maarel, E., 1989, 'Twenty years of numerical syntaxonomy', *Vegetation* 81, 1–15.
- Mueller-Dombois, D. & Ellenberg, H., 1974, *Aims and methods of vegetation ecology*, John Wiley and Sons, Inc., New York.
- O'Brien, J.W., 2004, 'Vegetation classification, mapping and condition assessment of Shamwari Game Reserve, Eastern Cape', MSc thesis, Terrestrial Ecology Research Unit, University of Port Elizabeth.
- Steenkamp, C., 1999, 'The Makuleke land claim power relations and CBNRM', draft report to the Evaluating Eden Project, International Institute for Environment and Development, London.